# **Dosing for Infants and Children**





From Your Healthcare Professional

#### DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.

TYLENOL If possible, use weight to dose: otherwise, use age. mL = milliliter

### Infants' TYLENOL® **Oral Suspension**

Active Ingredient: Acetaminophen 160 mg (in each 5 mL)

Available in: Grape

TYLENOL

### Children's TYLENOL® **Oral Suspension**

**Active Ingredient:** Acetaminophen 160 mg (in each 5 mL)

Use product only as directed



Use product only as directed

WEIGHT	AGE	Cherry	Cherry Use product only as directed.
6-11 POUNDS	0-3 MONTHS	1.25 mL	_
<b>12-17</b> POUNDS	4-11 MONTHS	2.5 mL	_
<b>18-23</b> POUNDS	12-23 MONTHS	3.75 mL	_
<b>24-35</b> POUNDS	2-3 YEARS	5 mL	5 mL 5 mL
36-47 POUNDS	4-5 YEARS	_	7.5 mL
<b>48-59</b> POUNDS	6-8 YEARS	_	10 mL
60-71 POUNDS	9-10 YEARS	_	12.5 mL
<b>72-95</b> POUNDS	11 YEARS	_	15 mL

### Kids' Wellness Tracker

This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App Store<sup>™</sup> and the Google Play Store™





Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

# **IMPORTANT INSTRUCTIONS** for Proper Use

Today's Date: \_

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all TYLENOL® products.
- Repeat dose every 4 hours while symptoms last.
- Do NOT give more than 5 doses in 24 hours.
- Do NOT use with any other product containing acetaminophen.
- Use only the dosing device (syringe or dosing cup) that came with the product.
- All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL).

# **Dosing for Infants and Children**

From Your Healthcare Professional





#### DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

Available in:

Motrin If possible, use weight to dose; otherwise, use age. Lasts up to 8 hours **mL** = milliliter

### Infants' MOTRIN® **Concentrated Drops**

**Active Ingredient:** 

Ibuprofen 50 mg (NSAID)\* (in each 1.25 mL)

\*Nonsteroidal



## Children's MOTRIN® **Oral Suspension**

**Active Ingredient:** Ibuprofen 100 mg (NSAID)\*

(in each 5 mL)

\*Nonsteroidal





			anti-inflammatory drug Berry		anti-inflammatory drug	Berry
WEIGHT	AGE	Serry Flavor Use pr	roduct only as directed.  Dye-Fr Berry	e Berry Flavor	Use product only as directed.	Dye-Free Berry
_	<6 MONTHS	<u> </u>		_		
<b>12-17</b> POUNDS	6-11 MONTHS		.825 mL 1.25 mL	_		
<b>18-23</b> POUNDS	12-23 MONTHS		6525 mL 1125 mL	_		
<b>24-35</b> POUNDS	2-3 YEARS			5 mL	5 mL	
<b>36-47</b> POUNDS	4-5 YEARS			7.5 ml	7.5 mL	
<b>48-59</b> POUNDS	6-8 YEARS			10 mL	10 mL	
60-71 POUNDS	9-10 YEARS			12.5 n	<b>nL</b> 12.5 mL	
<b>72-95</b> POUNDS	11 YEARS	-		15 mL	15 mL	

### Kids' Wellness Tracker

This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App Store<sup>™</sup> and the Google Play Store™





Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

# **IMPORTANT INSTRUCTIONS** for Proper Use

Today's Date: \_

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products.
- Repeat dose every 6-8 hours as needed.
- Do NOT give more than 4 doses in 24 hours.
- Use only the dosing device (syringe or dosing cup) that came with the product.